## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

#### (19) World Intellectual Property Organization International Bureau



# ) (1881 | 1881 | 1881 | 1881 | 1881 | 1881 | 1881 | 1881 | 1881 | 1881 | 1881 | 1881 | 1881 | 1881 | 1881 | 1881

#### (43) International Publication Date 18 October 2001 (18.10.2001)

#### PCT

### (10) International Publication Number WO 01/76661 A1

(51) International Patent Classification7: A61M 1/34, 1/16

[IT/IT]; Via S. Martino Carano, 84, I-41037 Mirandola

(21) International Application Number: PCT/IB01/00544

(74) Agent: LEJEUNE, Daniel; Hospal Service Brevets, 61, avenue Tony Garnier, F-69007 Lyon (FR).

(84) Designated States (regional): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,

(22) International Filing Date: 2 April 2001 (02.04.2001)

(25) Filing Language:

English

(81) Designated States (national): AU, CA, US.

(26) Publication Language:

TO2000A000333

English

(30) Priority Data:

7 April 2000 (07.04.2000)

(71) Applicant (for all designated States except US): HOSPAL AG [CH/CH]; Dornacherstrasse 8, CH-4008 Basel (CH).

Published: with international search report


NL, PT, SE, TR).

(72) Inventor; and

(75) Inventor/Applicant (for US only): PEDRAZZI, Renato

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DEVICE AND METHOD FOR CONTROLLING INFUSION OF A LIQUID IN AN EXTRACORPOREAL BLOOD **CIRCUIT** 



(57) Abstract: A Dialysis machine comprises: an extracorporeal blood circuit having an arterial pipe (12) and a venous pipe (15) respectively connected to an inlet and an outlet of a blood compartment (5) of a filter (4) having a blood compartment (5) and a dialysis liquid compartment (6) separated by a semi-permeable membrane (7); means (31) for determining a sequence of infusion of a liquid in the extracorporeal blood circuit, from at least one characteristic value (FF, TMP<sub>sve</sub>, K<sub>ef</sub>) correlated with the concentration of the blood (CE and/or a filtration efficiency of the filter (4); and means for infusing the infusion solution in the arterial pipe (12) and in the venous pipe (15) in accordance with the determined infusion sequence.